

Polynomials Related to Harmonic Numbers and Evaluation of Harmonic Number Series II

Ayhan Dil and Veli Kurt

Department of Mathematics, Akdeniz University, 07058 Antalya Turkey
adil@akdeniz.edu.tr, vkurt@akdeniz.edu.tr

February 3, 2010

Abstract

In this paper we focus on r -geometric polynomials, r -exponential polynomials and their harmonic versions. It is shown that harmonic versions of these polynomials and their generalizations are useful to obtain closed forms of some series related to harmonic numbers.

2000 Mathematics Subject Classification. 11B73, 11B75, 11B83

Key words: Exponential numbers and polynomials, Bell numbers and polynomials, geometric numbers and polynomials, Fubini numbers and polynomials, harmonic and hyperharmonic numbers.

1 Introduction

In [15] the concept of *harmonic*-geometric polynomials and *harmonic*-exponential polynomials are introduced and *hyperharmonic* generalizations of these polynomials and numbers are obtained. Furthermore it is shown that these polynomials are quite useful to obtain closed forms of some series related to harmonic numbers. In this paper, we extend this analysis to r -versions of these polynomials and numbers.

Boyadzhiev [6] has presented and discussed the following transformation formula:

$$\sum_{n=0}^{\infty} \frac{g^{(n)}(0)}{n!} f(n) x^n = \sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k g^{(k)}(x) \quad (1)$$

where f, g are appropriate functions and $\left\{ \begin{matrix} n \\ k \end{matrix} \right\}$ are Stirling numbers of the second kind.

One of the principal objectives of the present paper is to give closed forms of some series related to harmonic numbers as well. To this end, we give a useful generalization of (1) which contains r -Stirling numbers of the second kind instead of Stirling numbers of the second kind as:

$$\sum_{n=r}^{\infty} \frac{g^{(n)}(0)}{n!} \binom{n}{r} \frac{r!}{n^r} f_r(n) x^n = \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r x^k g^{(k)}(x), \quad (2)$$

where $f_r(x)$ denotes the Maclaurin series of $f(x)$ exclude the first $r - 1$ terms.

Thanks to formula (2) we introduce the concept of r -geometric and r -exponential polynomials and numbers. We obtain explicit relations between the r -versions and the classical versions of these polynomials and numbers. Besides, we present harmonic (and hyperharmonic) versions of r -geometric and r -exponential polynomials and numbers as well. The short lists of all these polynomials and numbers are given.

On the other hand formula (2) and harmonic r -geometric polynomials enables us to obtain closed forms of the following series

$$\sum_{n=r}^{\infty} \binom{n}{r} r! n^{m-r} H_n x^n, \quad (3)$$

where m and r are integers such that $m \geq r$ and H_n is the n -th partial sum of the harmonic series

In the rest of this section we introduce some important notions.

Stirling numbers of the first and second kind

Stirling numbers of the first kind $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]$ and Stirling numbers of the second kind $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ are quite important in combinatorics ([4, 5, 11, 21]). Briefly for the integers $n \geq k \geq 0$; $\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]$ counts the number of permutations of n elements with exactly k cycles and $\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}$ counts the number of ways to partition a set with n elements into k disjoint, nonempty subsets ([11]).

We note that for $n \geq k \geq 1$, the following identity holds for Stirling numbers of the second kind

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\} = \left\{ \begin{smallmatrix} n-1 \\ k-1 \end{smallmatrix} \right\} + k \left\{ \begin{smallmatrix} n-1 \\ k \end{smallmatrix} \right\}. \quad (4)$$

There is a certain generalization of these numbers namely r -Stirling numbers ([8]) which are similar to the weighted Stirling numbers ([9, 10]). Representation and combinatorial meanings of these numbers are as follows ([8]):

r -Stirling numbers of the first kind;

$$\left[\begin{smallmatrix} n \\ k \end{smallmatrix} \right]_r = \text{The number of permutations of the set } \{1, 2, \dots, n\} \text{ having} \\ k \text{ cycles, such that the numbers } 1, 2, \dots, r \text{ are in distinct cycles,}$$

r -Stirling numbers of the second kind;

$$\left\{ \begin{smallmatrix} n \\ k \end{smallmatrix} \right\}_r = \text{The number of partitions of the set } \{1, 2, \dots, n\} \text{ into} \\ k \text{ non-empty disjoint subsets, such that the numbers} \\ 1, 2, \dots, r \text{ are in distinct subsets.}$$

Specializing $r = 0$ gives the classical Stirling numbers.

The r -Stirling numbers of the second kind satisfy the same recurrence relation as (4) except for the initial conditions, i.e ([8])

$$\begin{aligned}\left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r &= 0, \quad n < r, \\ \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r &= \delta_{k,r}, \quad n = r, \\ \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r &= \left\{ \begin{matrix} n-1 \\ k-1 \end{matrix} \right\}_r + k \left\{ \begin{matrix} n-1 \\ k \end{matrix} \right\}_r, \quad n > r.\end{aligned}\tag{5}$$

Exponential polynomials and numbers

Exponential polynomials (or single variable Bell polynomials) $\phi_n(x)$ are defined by([2, 7, 16, 21])

$$\phi_n(x) := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} x^k.\tag{6}$$

The first few exponential polynomials are:

$\phi_0(x) = 1,$	(7)
$\phi_1(x) = x,$	
$\phi_2(x) = x + x^2,$	
$\phi_3(x) = x + 3x^2 + x^3,$	
$\phi_4(x) = x + 7x^2 + 6x^3 + x^4.$	

The well known exponential numbers (or Bell numbers) are obtained by setting $x = 1$ in $\phi_n(x)$, i.e ([3, 11, 12])

$$\phi_n := \phi_n(1) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}.\tag{8}$$

Hence the first few exponential numbers are:

$$\phi_0 = 1, \phi_1 = 1, \phi_2 = 2, \phi_3 = 5, \phi_4 = 15.\tag{9}$$

In [14] the authors obtained some fundamental properties of the exponential polynomials and numbers using Euler-Seidel matrices method as:

$$\phi_{n+1}(x) = x \sum_{k=0}^n \binom{n}{k} \phi_k(x)\tag{10}$$

and

$$\phi_{n+1} = \sum_{k=0}^n \binom{n}{k} \phi_k.\tag{11}$$

Recently, Mezo ([18]) has defined the " r -Bell polynomials and numbers" as:

$$B_{n,r}(x) = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r x^k\tag{12}$$

and

$$B_{n,r} = \sum_{k=0}^n \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r \quad (13)$$

respectively. r -Exponential polynomials and numbers which we discuss in the present paper are slightly different than the r -Bell polynomials and numbers in [18].

Geometric polynomials and numbers

Geometric polynomials are defined in [6, 22, 23] as follows:

$$F_n(x) := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} k! x^k. \quad (14)$$

The first few geometric polynomials are:

$F_0(x) = 1,$	(15)
$F_1(x) = x,$	
$F_2(x) = x + 2x^2,$	
$F_3(x) = x + 6x^2 + 6x^3,$	
$F_4(x) = x + 14x^2 + 36x^3 + 24x^4.$	

Specializing $x = 1$ in (14) we get geometric numbers (or ordered Bell numbers) F_n as ([6, 23, 24]):

$$F_n := F_n(1) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} k!. \quad (16)$$

The first few geometric numbers are:

$$F_0 = 1, F_1 = 1, F_2 = 3, F_3 = 13, F_4 = 75. \quad (17)$$

Boyadzhiev ([6]) introduced the "general geometric polynomials" as

$$F_{n,r}(x) = \frac{1}{\Gamma(r)} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\} \Gamma(k+r) x^k, \quad (18)$$

where $\text{Re}(r) > 0$. In the fifth section we will deal with the general geometric polynomials.

Exponential and geometric polynomials are connected by the following integral relation ([6])

$$F_n(z) = \int_0^\infty \phi_n(z\lambda) e^{-\lambda} d\lambda. \quad (19)$$

In [14] the authors also obtained some fundamental properties of the geometric polynomials and numbers using Euler-Seidel matrices method as:

$$F_{n+1}(x) = x \sum_{k=0}^n \binom{n+1}{k} F_k(x) \quad (20)$$

and

$$F_n = \sum_{k=0}^{n-1} \binom{n}{k} F_k. \quad (21)$$

By means of r -Stirling numbers Nyul ([20]) introduced " r -geometric polynomials and numbers (or r -Fubini or ordered r -Bell polynomials and numbers)" are respectively as follows:

$$F_{n,r}(x) = \sum_{k=0}^n (k+r)! \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r x^k \quad (22)$$

and

$$F_{n,r} = \sum_{k=0}^n (k+r)! \left\{ \begin{matrix} n+r \\ k+r \end{matrix} \right\}_r.$$

In this work r -geometric polynomials come up naturally in an application of the generalized transformation formula as well.

Notice that, our concept of r -geometric polynomials is slightly different than in [20].

Harmonic and Hyperharmonic numbers

The n -th harmonic number is the n -th partial sum of the harmonic series:

$$H_n := \sum_{k=1}^n \frac{1}{k}, \quad (23)$$

where $H_0 = 0$.

For an integer $\alpha > 1$, let

$$H_n^{(\alpha)} := \sum_{k=1}^n H_k^{(\alpha-1)}, \quad (24)$$

with $H_n^{(1)} := H_n$, be the n -th hyperharmonic number of order α ([4, 12]).

These numbers can be expressed in terms of binomial coefficients and ordinary harmonic numbers as ([12, 19]):

$$H_n^{(\alpha)} = \binom{n+\alpha-1}{\alpha-1} (H_{n+\alpha-1} - H_{\alpha-1}). \quad (25)$$

Well-known generating functions of the harmonic and hyperharmonic numbers are given by

$$\sum_{n=1}^{\infty} H_n x^n = -\frac{\ln(1-x)}{1-x} \quad (26)$$

and

$$\sum_{n=1}^{\infty} H_n^{(\alpha)} x^n = -\frac{\ln(1-x)}{(1-x)^\alpha}. \quad (27)$$

respectively ([13]).

The following relations connect harmonic and hyperharmonic numbers with the Stirling and r -Stirling numbers of the first kind ([4]):

$$\begin{bmatrix} k+1 \\ 2 \end{bmatrix} = k!H_k, \quad (28)$$

and

$$k!H_k^{(r)} = \begin{bmatrix} n+r \\ r+1 \end{bmatrix}_r. \quad (29)$$

2 Generalization of the transformation formula

In this section firstly we mention Boyadzhiev's Theorem 4.1 in [6] and give a useful generalization of it. As a result of this generalization we introduce r -geometric polynomials and numbers.

Suppose we are given an entire function f and a function g , analytic in a region containing the annulus $K = \{z : r < |z| < R\}$ where $0 < r < R$. Hence these functions have following series expansions,

$$f(x) = \sum_{n=0}^{\infty} p_n x^n \text{ and } g(x) = \sum_{n=-\infty}^{\infty} q_n x^n.$$

Now we are ready to state Boyadzhiev's theorem.

Theorem 1 ([6]) *Let the functions f and g be described as above. If the series*

$$\sum_{n=-\infty}^{\infty} q_n f(n) x^n$$

converges absolutely on K , then

$$\sum_{n=-\infty}^{\infty} q_n f(n) x^n = \sum_{m=0}^{\infty} p_m \sum_{k=0}^m \begin{Bmatrix} m \\ k \end{Bmatrix} x^k g^{(k)}(x) \quad (30)$$

holds for all $x \in K$.

Stirling numbers of the second kind appear in the formula (30) due to (xD) operator. Our aim is to get a more general formula than (30) which contains r -Stirling numbers of the second kind instead of Stirling numbers of the second kind. Accordingly, first we generalize the operator (xD) .

2.1 Generalization of the operator (xD)

The operator (xD) operates a function $f(x)$ as;

$$(xD)f(x) := xf'(x), \quad (31)$$

where f' is the first derivative of the function f .

For any m -times differentiable function f we have ([6]),

$$(xD)^m f(x) = \sum_{k=0}^m \left\{ \begin{matrix} m \\ k \end{matrix} \right\} x^k f^{(k)}(x). \quad (32)$$

This fact can be easily proven with induction on m by the help of (4).

Our first aim is to generalize the operator (xD) . Later we use this generalization to obtain r -geometric and r -exponential polynomials and numbers. In the light of this motivation and after a plenty of observations we arrive the following definition.

Definition 2 *Let f be a function which is at least m -times differentiable and r be a nonnegative integer. Then the action (xD_r) is*

$$(xD_r)^m f(x) := \begin{cases} 0 & , m < r \\ (xD)^{m-r} x^r f^{(r)}(x) & , m \geq r \end{cases}. \quad (33)$$

An equivalent statement of this definition is given by the following proposition.

Proposition 3 *By applying (xD_r) m -times to a function f which is at least m -times differentiable, then the following equation holds*

$$(xD_r)^m f(x) = \sum_{k=0}^m \left\{ \begin{matrix} m \\ k \end{matrix} \right\}_r x^k f^{(k)}(x), \quad (34)$$

where $m \geq r$.

Proof. It follows from induction on m , in the light of Definition 2 and recurrence relation (5). ■

The equation (34) is a generalization of the equation (32) since setting $r = 0$ in (34) gives the equation (32).

Corollary 4 *Let n be an integer, then*

$$(xD_r)^m x^n = n^{m-r} \binom{n}{r} r! x^n. \quad (35)$$

2.2 Generalization of the transformation formula

Now we give our main theorem that is a generalization of Theorem 1.

Theorem 5 *Let $f(x)$ be an entire function and $g(x)$ be an analytic function on the annulus $K = \{z, s < |z| < S\}$, where $0 \leq s < S$. Suppose that their power series be given as*

$$f(x) = \sum_{m=0}^{\infty} p_m x^m \text{ and } g(x) = \sum_{n=-\infty}^{\infty} q_n x^n.$$

If the series

$$\sum_{n=-\infty}^{\infty} q_n \binom{n}{r} \frac{r!}{n^r} f_r(n) x^n \quad (36)$$

where r is a nonnegative integer and $f_r(x)$ denotes the power series $\sum_{m=r}^{\infty} p_m x^m$, converges absolutely on K , then

$$\sum_{n=-\infty}^{\infty} q_n \binom{n}{r} \frac{r!}{n^r} f_r(n) x^n = \sum_{m=r}^{\infty} p_m \sum_{k=0}^m \left\{ \begin{matrix} m \\ k \end{matrix} \right\}_r x^k g^{(k)}(x) \quad (37)$$

holds for all $x \in K$.

Proof. By considering the power series expansion of $g(x)$ with (34) and (35) we have

$$\sum_{n=-\infty}^{\infty} q_n \binom{n}{r} n^{m-r} r! x^n = \sum_{k=0}^m \left\{ \begin{matrix} m \\ k \end{matrix} \right\}_r x^k g^{(k)}(x) \quad (38)$$

where m and r are integer such that $m \geq r \geq 0$. If we multiply both sides of the equation (38) by p_m and sum on m from r to infinity we get

$$\sum_{n=-\infty}^{\infty} q_n \binom{n}{r} \frac{r!}{n^r} \sum_{m=r}^{\infty} p_m n^m x^n = \sum_{m=r}^{\infty} p_m \sum_{k=0}^m \left\{ \begin{matrix} m \\ k \end{matrix} \right\}_r x^k g^{(k)}(x),$$

since (36) is converges absolutely on K . This completes proof. ■

Corollary 6 Let g be an analytic function on the disk $D = \{z, 0 \leq |z| < S\}$ then

$$\sum_{n=r}^{\infty} \frac{g^{(n)}(0)}{n!} \binom{n}{r} \frac{r!}{n^r} f_r(n) x^n = \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r x^k g^{(k)}(x). \quad (39)$$

Most of the results in the subsequent sections depend on the Corollary 6.

Remark 7 Specializing $r = 0$ in the Theorem 5 we turn back to the Theorem 4.1 of Boyadzhiev ([6]). Therefore from now on we are interested in the case $r \geq 1$.

3 r -exponential and r -geometric polynomials and numbers

Stirling numbers of the first and second kind are notable in many branches of mathematics, especially in combinatorics, computational mathematics and computer sciences ([1, 5, 11, 12, 17]). Importance of the exponential polynomials and numbers are substantially because of their direct connection with Stirling numbers. r -Stirling numbers ([8]) are one of the reputable generalizations of Stirling numbers. Therefore introduction of the concepts of the r -exponential and r -geometric polynomials and numbers are good motivation for us.

3.1 r -exponential polynomials and numbers

Firstly we consider $g(x) = e^x$ in the equation (39). Hence we get

$$\sum_{n=r}^{\infty} \binom{n}{r} \frac{r!}{n^r} f_r(n) \frac{x^n}{n!} = e^x \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r x^k. \quad (40)$$

The finite sum on the RHS is a generalization of exponential polynomials. We call these polynomials as " r -exponential polynomials" and indicate them with ${}_r\phi_n(x)$. Hence

$${}_r\phi_n(x) := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r x^k. \quad (41)$$

The first few r -exponential polynomials are:

${}_r\phi_n(x)$	$r = 1$	$r = 2$	$r = 3$
$n = 0$	0	0	0
$n = 1$	x	0	0
$n = 2$	$x + x^2$	x^2	0
$n = 3$	$x + 3x^2 + x^3$	$2x^2 + x^3$	x^3
$n = 4$	$x + 7x^2 + 6x^3 + x^4$	$4x^2 + 5x^3 + x^4$	$3x^3 + x^4$

(42)

Similar to the classical case, " r -exponential numbers" can be defined by setting $x = 1$ in (41) i.e,

$${}_r\phi_n := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r. \quad (43)$$

Hence the first few r -exponential numbers are:

${}_r\phi_n$	$r = 1$	$r = 2$	$r = 3$
$n = 0$	0	0	0
$n = 1$	1	0	0
$n = 2$	2	1	0
$n = 3$	5	3	1
$n = 4$	15	10	4

(44)

Now we give an explicit formula which connects r -exponential polynomials with the classical exponential polynomials. Also this formula allows us to calculate ${}_r\phi_n(x)$ easily.

Proposition 8 *We have*

$${}_r\phi_{n+r}(x) = x^r \sum_{k=0}^n \binom{n}{k} r^{n-k} \phi_k(x), \quad (45)$$

where n and r are nonnegative integers.

Proof. Let m be an integer such that $m \geq r \geq 0$ and we specialize $f(x) = x^m$ in (40). Then we get

$${}_r\phi_m(x) e^x = \sum_{n=r}^{\infty} \binom{n}{r} \frac{r!}{n!} n^{m-r} x^n.$$

RHS of this equation can be written as

$$x^r \sum_{n=0}^{\infty} (n+r)^{m-r} \frac{x^n}{n!} = x^r \sum_{k=0}^{m-r} \binom{m-r}{k} r^{m-r-k} \sum_{n=0}^{\infty} n^k \frac{x^n}{n!}.$$

Considering the definition of the operator (xD) this becomes

$$x^r \sum_{k=0}^{m-r} \binom{m-r}{k} r^{m-r-k} (xD)^k e^x.$$

The equation (32) enables us to write

$$x^r e^x \sum_{k=0}^{m-r} \binom{m-r}{k} r^{m-r-k} \phi_k(x).$$

Comparison of the LHS and the RHS completes the proof. ■

Similar relation can be given between classical exponential numbers and r -exponential numbers as a corollary.

Corollary 9

$${}_r\phi_{n+r} = \sum_{k=0}^n \binom{n}{k} r^{n-k} \phi_k. \quad (46)$$

The following corollary shows that the equation (45) is a generalization of the equation (10).

Corollary 10

$$\phi_{n+1}(x) = x \sum_{k=0}^n \binom{n}{k} \phi_k(x).$$

3.2 r -geometric polynomials and numbers

By considering $g(x) = \frac{1}{1-x}$ in the equation (39) we get

$$\sum_{n=r}^{\infty} \binom{n}{r} \frac{r!}{n^r} f_r(n) x^n = \frac{1}{1-x} \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! \left(\frac{x}{1-x} \right)^k. \quad (47)$$

We call the finite sum of the RHS as " r -geometric polynomials" and indicate them with ${}_rF_n(x)$. Hence

$${}_rF_n(x) := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! x^k. \quad (48)$$

The first few r -geometric polynomials are:

${}_rF_n(x)$	$r = 1$	$r = 2$	$r = 3$
$n = 0$	0	0	0
$n = 1$	x	0	0
$n = 2$	$x + 2x^2$	$2x^2$	0
$n = 3$	$x + 6x^2 + 6x^3$	$4x^2 + 6x^3$	$6x^3$
$n = 4$	$x + 14x^2 + 36x^3 + 24x^4$	$8x^2 + 30x^3 + 24x^4$	$18x^3 + 24x^4$

(49)

We define " r -geometric numbers" by specializing $x = 1$ in (48) as

$${}_rF_n := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k!. \quad (50)$$

The first few r -geometric numbers are:

${}_rF_n$	$r = 1$	$r = 2$	$r = 3$
$n = 0$	0	0	0
$n = 1$	1	0	0
$n = 2$	3	2	0
$n = 3$	13	10	6
$n = 4$	75	62	42

(51)

The following proposition gives an explicit formula between r -geometric polynomials and generalized geometric polynomials which have given by the equation (18).

Proposition 11 *For any nonnegative integers n and r we have*

$${}_rF_{n+r}(x) = x^r r! \sum_{k=0}^n \binom{n}{k} r^{n-k} F_{k,r+1}(x) \quad (52)$$

Proof. Let m be a nonnegative integer such that $m \geq r$. By setting $f(x) = x^m$ in (47) we get

$$\frac{1}{1-x} {}_rF_m\left(\frac{x}{1-x}\right) = \sum_{n=r}^{\infty} \binom{n}{r} r! n^{m-r} x^n.$$

Rearranging RHS gives

$$x^r r! \sum_{k=0}^{m-r} \binom{m-r}{k} r^{m-r-k} \sum_{n=0}^{\infty} \binom{n+r}{r} n^k x^n.$$

We can write this by means of (xD) operator as

$$x^r r! \sum_{k=0}^{m-r} \binom{m-r}{k} r^{m-r-k} (xD)^k \frac{1}{(1-x)^{r+1}}.$$

Considering the fact that (equation (3.26) in [6])

$$(xD)^k \frac{1}{(1-x)^{r+1}} = \frac{1}{(1-x)^{r+1}} F_{k,r+1} \left(\frac{x}{1-x} \right)$$

completes the proof. ■

A similar result between numbers is as follows.

Corollary 12

$${}_r F_{n+r} = r! \sum_{k=0}^n \binom{n}{k} r^{n-k} F_{k,r+1} \quad (53)$$

Owing to (53), we give the following relations for classical geometric polynomials and numbers as a corollary.

Corollary 13

$$F_{n+1}(x) = x \sum_{k=0}^n \binom{n}{k} F_{k,2}(x), \quad (54)$$

$$F_{n+1} = \sum_{k=0}^n \binom{n}{k} F_{k,2}. \quad (55)$$

4 Harmonic r -geometric and harmonic r -exponential polynomials and numbers

We introduce the concepts of *harmonic*-geometric and *harmonic*-exponential polynomials and numbers in [15]. Along this section we follow similar approach in ([15]) to investigate harmonic r -geometric and harmonic r -exponential polynomials and numbers.

4.1 Harmonic r -geometric polynomials and numbers

We consider the generating function of harmonic numbers as the function g in the transformation formula (39). From [15] we have

$$g^{(k)}(z) = \frac{k! (H_k - \ln(1-z))}{(1-z)^{k+1}} \quad (56)$$

and

$$g^{(k)}(0) = k! H_k. \quad (57)$$

With the help of Theorem 5 we state the following transformation formula for harmonic numbers.

Proposition 14 *Let r be a nonnegative integer and f be an entire function. Then we have*

$$\begin{aligned} & \sum_{n=r}^{\infty} \binom{n}{r} H_n \frac{r!}{n^r} f_r(n) x^n \\ = & \frac{1}{1-x} \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! H_k \left(\frac{x}{1-x} \right)^k \\ & - \frac{\ln(1-x)}{1-x} \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! \left(\frac{x}{1-x} \right)^k. \end{aligned} \quad (58)$$

Proof. Employing (56) and (57) in (39) gives the statement. ■

Second part of the RHS of the equation (58) contains r -geometric polynomials which are familiar to us from the previous section. But the first part contains a new family of polynomials which is a generalization of *harmonic*-geometric polynomials ([15]). We call them as "harmonic r -geometric polynomials and indicate them with ${}_r F_n^h(x)$. Thus

$${}_r F_n^h(x) := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! H_k x^k. \quad (59)$$

The first few harmonic r -geometric polynomials are:

${}_r F_n^h(x)$	$r = 1$	$r = 2$	$r = 3$
$n = 0$	0	0	0
$n = 1$	x	0	0
$n = 2$	$x + 3x^2$	$3x^2$	0
$n = 3$	$x + 9x^2 + 11x^3$	$6x^2 + 11x^3$	$11x^3$
$n = 4$	$x + 21x^2 + 66x^3 + 50x^4$	$12x^2 + 55x^3 + 50x^4$	$33x^3 + 50x^4$

(60)

"Harmonic r -geometric numbers" can be defined by setting $x = 1$ in (59), i.e

$${}_r F_n^h := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! H_k. \quad (61)$$

The first few harmonic r -geometric numbers are:

${}_r F_n^h$	$r = 1$	$r = 2$	$r = 3$
$n = 0$	0	0	0
$n = 1$	1	0	0
$n = 2$	4	3	0
$n = 3$	21	17	11
$n = 4$	138	117	83

(62)

Hence with this notation we state the formula (58) simply as

$$\begin{aligned} & \sum_{n=r}^{\infty} \binom{n}{r} H_n \frac{r!}{n^r} f_r(n) x^n \\ &= \frac{1}{1-x} \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \left\{ {}_r F_n^h \left(\frac{x}{1-x} \right) - {}_r F_n \left(\frac{x}{1-x} \right) \ln(1-x) \right\}. \end{aligned} \quad (63)$$

Due to the following corollary we obtain closed forms of some series related to harmonic numbers and binomial coefficients.

Corollary 15

$$\sum_{n=r}^{\infty} \binom{n}{r} r! n^{m-r} H_n x^n = \frac{1}{1-x} \left\{ {}_r F_m^h \left(\frac{x}{1-x} \right) - {}_r F_m \left(\frac{x}{1-x} \right) \ln(1-x) \right\}, \quad (64)$$

where m and r are integers such that $m \geq r$.

Proof. It follows by setting $f(x) = x^m$ in the equation (63). ■

Remark 16 Formula (64) allow us to calculate closed forms of several harmonic number series. The case $r = 1$ in (64) has been analyzed in [15] already.

The case $r = 2$ gives

$$\sum_{n=2}^{\infty} n^{m-1} (n-1) H_n x^n = \frac{1}{1-x} \left\{ {}_2 F_m^h \left(\frac{x}{1-x} \right) - {}_2 F_m \left(\frac{x}{1-x} \right) \ln(1-x) \right\}. \quad (65)$$

Hence some series and their closed forms that we get from (65) are as follows:

For $m = 2$ we have

$$\sum_{n=2}^{\infty} n(n-1) H_n x^n = \frac{x^2 \{3 - 2 \ln(1-x)\}}{(1-x)^3}. \quad (66)$$

For $m = 3$ we have

$$\sum_{n=2}^{\infty} n^2 (n-1) H_n x^n = \frac{x^2 \{6 + 5x - (4 + 2x) \ln(1-x)\}}{(1-x)^4}, \quad (67)$$

and so on.

The case $r = 3$ gives

$$\begin{aligned} & \sum_{n=3}^{\infty} n^{m-2} (n-1)(n-2) H_n x^n \\ &= \frac{1}{1-x} \left\{ {}_3 F_m^h \left(\frac{x}{1-x} \right) - {}_3 F_m \left(\frac{x}{1-x} \right) \ln(1-x) \right\}. \end{aligned} \quad (68)$$

Hence some series and their closed forms that we get from (68) are as follows:

For $m = 3$ we have

$$\sum_{n=3}^{\infty} n(n-1)(n-2)H_n x^n = \frac{x^3 \{11 - 6 \ln(1-x)\}}{(1-x)^4}. \quad (69)$$

For $m = 4$ we have

$$\sum_{n=3}^{\infty} n^2(n-1)(n-2)H_n x^n = \frac{x^3 \{33 + 17x - (18 + 6x) \ln(1-x)\}}{(1-x)^5}, \quad (70)$$

and so on.

Now we give a summation formula for the multiple series.

Proposition 17

$$\begin{aligned} & \sum_{n=r}^{\infty} \left(\sum_{k=0}^{n-r} \binom{k}{r} \binom{n+s-k}{s} r! k^{m-r} H_k \right) x^n \\ &= \sum_{n=r}^{\infty} \left(\sum_{0 \leq k_1 \leq k_2 \leq \dots \leq k_{s+1} \leq n} \binom{k_1}{r} r! k_1^{m-r} H_{k_1} \right) x^n \\ &= \frac{1}{(1-x)^{s+2}} \left\{ {}_r F_m^h \left(\frac{x}{1-x} \right) - {}_r F_m \left(\frac{x}{1-x} \right) \ln(1-x) \right\} \end{aligned} \quad (71)$$

Proof. Multiplying both sides of the equation (64) with the Newton binomial series and considering that

$$\sum_{k=0}^{n-r} \binom{k}{r} \binom{n+s-k}{s} r! k^{m-r} H_k = \sum_{0 \leq k_1 \leq k_2 \leq \dots \leq k_{s+1} \leq n} \binom{k_1}{r} r! k_1^{m-r} H_{k_1}$$

we get the statement. ■

By setting $r = 2$ and $s = 0$ in the formula (71) we can give the following applications:

For $m = 2$ we have

$$\sum_{n=2}^{\infty} \left(\sum_{k=2}^n k(k-1)H_k \right) x^n = \frac{x^2 \{3 - 2 \ln(1-x)\}}{(1-x)^4}. \quad (72)$$

For $m = 3$ we have

$$\sum_{n=2}^{\infty} \left(\sum_{k=2}^n k^2(k-1)H_k \right) x^n = \frac{x^2 \{6 + 5x - (4 + 2x) \ln(1-x)\}}{(1-x)^5}, \quad (73)$$

and so on.

Remark 18 By the help of (28) we can state ${}_rF_n^h(x)$ and ${}_rF_n^h$ in terms of r -Stirling numbers of the second kind and Stirling numbers of the first kind as

$${}_rF_n^h(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r \begin{bmatrix} k+1 \\ 2 \end{bmatrix} x^k \quad (74)$$

and

$${}_rF_n^h = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r \begin{bmatrix} k+1 \\ 2 \end{bmatrix}. \quad (75)$$

4.2 Harmonic r -exponential polynomials and numbers

Bearing in mind the similarity of exponential and geometric polynomials and being inspired by the definition of harmonic exponential polynomials and numbers we arrive the following definition.

Definition 19 For the nonnegative integers n and r , "harmonic r -exponential polynomials" and "harmonic r -exponential numbers" are defined respectively as

$${}_r\phi_n^h(x) := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r H_k x^k \quad (76)$$

and

$${}_r\phi_n^h := \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r H_k. \quad (77)$$

The first few harmonic r -exponential polynomials are:

${}_r\phi_n^h(x)$	$r = 1$	$r = 2$	$r = 3$
$n = 0$	0	0	0
$n = 1$	x	0	0
$n = 2$	$x + \frac{3}{2}x^2$	$\frac{3}{2}x^2$	0
$n = 3$	$x + \frac{9}{2}x^2 + \frac{11}{6}x^3$	$3x^2 + \frac{11}{6}x^3$	$\frac{11}{6}x^3$
$n = 4$	$x + \frac{21}{2}x^2 + 11x^3 + \frac{25}{12}x^4$	$6x^2 + \frac{55}{6}x^3 + \frac{25}{12}x^4$	$\frac{11}{2}x^3 + \frac{25}{12}x^4$

(78)

The first few harmonic r -exponential numbers are:

${}_r\phi_n^h$	$r = 1$	$r = 2$	$r = 3$
$n = 0$	0	0	0
$n = 1$	1	0	0
$n = 2$	$\frac{3}{2}$	$\frac{3}{2}$	0
$n = 3$	$\frac{22}{3}$	$\frac{29}{6}$	$\frac{11}{6}$
$n = 4$	$\frac{295}{12}$	$\frac{69}{4}$	$\frac{91}{12}$

(79)

Remark 20 Definition 19 enables us to extend the relation (19) as

$${}_rF_n^h(z) = \int_0^\infty {}_r\phi_n^h(z\lambda) e^{-\lambda} d\lambda. \quad (80)$$

5 Hyperharmonic r -geometric and hyperharmonic r -exponential polynomials and numbers

For the completeness of this work, now we consider hyperharmonic numbers and their transformations. In this way we could generalize almost all results of [15] and in previous sections of the present paper.

5.1 Hyperharmonic r -geometric polynomials and numbers

Similar to the previous section, let us consider the function g in the transformation formula (39) as the generating function of the hyperharmonic numbers. From [15] we have

$$g^{(k)}(x) = \frac{\Gamma(k+\alpha)}{\Gamma(\alpha)} \frac{1}{(1-z)^{\alpha+k}} (H_{k+\alpha-1} - H_{\alpha-1} - \ln(1-x)) \quad (81)$$

and

$$g^{(k)}(0) = k! H_k^{(\alpha)}. \quad (82)$$

Now we give a transformation formula for hyperharmonic numbers.

Proposition 21 *For integers $r \geq 0$ and $\alpha \geq 1$ we have*

$$\begin{aligned} & \sum_{n=r}^{\infty} \binom{n}{r} H_n^{(\alpha)} \frac{r!}{n^r} f_r(n) x^n \\ &= \frac{1}{(1-z)^\alpha} \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! H_k^{(\alpha)} \left(\frac{x}{1-x} \right)^k \\ & \quad - \frac{\ln(1-x)}{(1-z)^\alpha} \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \frac{1}{\Gamma(\alpha)} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r \Gamma(k+\alpha) \left(\frac{x}{1-x} \right)^k. \end{aligned} \quad (83)$$

Proof. Consideration (81) and (82) in (39) give the statement. ■

The first part of the RHS is a generalization of harmonic r -geometric polynomials which contains hyperharmonic numbers instead of harmonic numbers. We call these polynomials as "hyperharmonic r -geometric polynomials" and indicate them with ${}_r F_{n,\alpha}^h(x)$. Thus

$${}_r F_{n,\alpha}^h(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! H_k^{(\alpha)} x^k \quad (84)$$

The first few hyperharmonic r -geometric polynomials are:

Case $\alpha = 2$

${}_rF_{n,2}^h(x)$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	x	0
$n = 2$	$x + 5x^2$	$5x^2$
$n = 3$	$x + 15x^2 + 26x^3$	$10x^2 + 26x^3$
$n = 4$	$x + 35x^2 + 156x^3 + 154x^4$	$20x^2 + 130x^3 + 154x^4$

(85)

Case $\alpha = 3$

${}_rF_{n,3}^h(x)$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	x	0
$n = 2$	$x + 7x^2$	$7x^2$
$n = 3$	$x + 21x^2 + 47x^3$	$14x^2 + 47x^3$
$n = 4$	$x + 49x^2 + 282x^3 + 342x^4$	$28x^2 + 235x^3 + 342x^4$

(86)

The second part of the RHS of (83) contains also a generalization of the polynomials, "general geometric polynomials", which we mention with the equation (18). We call these polynomials as "general r -geometric polynomials" and indicate them with ${}_rF_{n,\alpha}(x)$. Hence

$${}_rF_{n,\alpha}(x) = \frac{1}{\Gamma(\alpha)} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r \Gamma(k + \alpha) x^k. \quad (87)$$

The first few general r -geometric polynomials are:

Case $\alpha = 2$

${}_rF_{n,2}(x)$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	$2x$	0
$n = 2$	$2x + 6x^2$	$6x^2$
$n = 3$	$2x + 18x^2 + 24x^3$	$12x^2 + 24x^3$
$n = 4$	$2x + 42x^2 + 144x^3 + 120x^4$	$24x^2 + 120x^3 + 120x^4$

(88)

and

Case $\alpha = 3$

${}_rF_{n,3}(x)$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	$3x$	0
$n = 2$	$3x + 12x^2$	$12x^2$
$n = 3$	$3x + 36x^2 + 60x^3$	$24x^2 + 60x^3$
$n = 4$	$3x + 84x^2 + 360x^3 + 360x^4$	$48x^2 + 300x^3 + 360x^4$

(89)

With the help of these notations we can state (83) simply as

$$\begin{aligned} & \sum_{n=r}^{\infty} \binom{n}{r} H_n^{(\alpha)} \frac{r!}{n^r} f(n) x^n \\ &= \frac{1}{(1-z)^\alpha} \sum_{n=r}^{\infty} \frac{f^{(n)}(0)}{n!} \left[{}_rF_{n,\alpha}^h \left(\frac{x}{1-x} \right) - {}_rF_{n,\alpha} \left(\frac{x}{1-x} \right) \ln(1-x) \right]. \end{aligned} \quad (90)$$

Remark 22 Specializing $x = 1$ in (84) we get "hyperharmonic r -geometric numbers" as

$${}_rF_{n,\alpha}^h = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r k! H_k^{(\alpha)}. \quad (91)$$

The first few hyperharmonic r -geometric numbers are:

Case $\alpha = 2$

${}_rF_{n,2}^h$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	1	0
$n = 2$	6	5
$n = 3$	42	36
$n = 4$	346	304

(92)

Case $\alpha = 3$

${}_rF_{n,3}^h$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	1	0
$n = 2$	8	7
$n = 3$	69	61
$n = 4$	674	605

(93)

and specializing $x = 1$ in (87) gives "general r -geometric numbers" as

$${}_rF_{n,\alpha} = \frac{1}{\Gamma(\alpha)} \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r \Gamma(k + \alpha). \quad (94)$$

The first few general r -geometric numbers are:

Case $\alpha = 2$

${}_rF_{n,2}$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	2	0
$n = 2$	8	6
$n = 3$	44	36
$n = 4$	308	264

(95)

and

Case $\alpha = 3$

${}_rF_{n,3}$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	3	0
$n = 2$	15	12
$n = 3$	99	84
$n = 4$	807	708

(96)

Thanks to the following corollary of Proposition (21) we have closed forms of some series related to hyperharmonic numbers and binomial coefficients.

Corollary 23

$$\sum_{n=r}^{\infty} \binom{n}{r} r! n^{m-r} H_n^{(\alpha)} x^n = \frac{1}{(1-z)^\alpha} \left[{}_rF_{m,\alpha}^h \left(\frac{x}{1-x} \right) - {}_rF_{m,\alpha} \left(\frac{x}{1-x} \right) \ln(1-x) \right]. \quad (97)$$

Proof. For a positive integers $m \geq r$, setting $f(x) = x^m$ in (90) gives (97). ■

Remark 24 Specializing the values of r , m and α in (97) one can get closed forms of several hyperharmonic numbers series.

Now we extend the formula (71) to hyperharmonic number series.

Proposition 25

$$\begin{aligned} & \sum_{n=r}^{\infty} \left(\sum_{k=0}^{n-r} \binom{k}{r} \binom{n+s-k}{s} r! k^{m-r} H_k^{(\alpha)} \right) x^n \\ &= \sum_{n=r}^{\infty} \left(\sum_{0 \leq k_1 \leq k_2 \leq \dots \leq k_{s+1} \leq n} \binom{k_1}{r} r! k_1^{m-r} H_{k_1}^{(\alpha)} \right) x^n \\ &= \frac{1}{(1-x)^{\alpha+s+1}} \left\{ {}_rF_m^h \left(\frac{x}{1-x} \right) - {}_rF_m \left(\frac{x}{1-x} \right) \ln(1-x) \right\} \end{aligned} \quad (98)$$

Proof. Multiplying both sides of the equation (97) with the Newton binomial series gives the statement. ■

Remark 26 Also special values of r , m , s and α in (98) gives closed forms of several mutiplicative hyperharmonic numbers series.

Remark 27 Using (29) we get an alternative expression of hyperharmonic r -geometric polynomials and numbers as

$${}_rF_{n,\alpha}^h(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r \left[\begin{matrix} k+\alpha \\ \alpha+1 \end{matrix} \right]_\alpha x^k, \quad (99)$$

$${}_rF_{n,\alpha}^h = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r \left[\begin{matrix} k+\alpha \\ \alpha+1 \end{matrix} \right]_\alpha. \quad (100)$$

5.2 Hyperharmonic r -exponential polynomials and numbers

Definition 28 For positive integers m and r "hyperharmonic r -exponential polynomials" are defined as

$${}_r\phi_{n,\alpha}^h(x) = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r H_k^{(\alpha)} x^k. \quad (101)$$

The first few hyperharmonic r -exponential polynomials are:
Case $\alpha = 2$

${}_r\phi_{n,2}^h(x)$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	x	0
$n = 2$	$x + \frac{5}{2}x^2$	$\frac{5}{2}x^2$
$n = 3$	$x + \frac{15}{2}x^2 + \frac{13}{3}x^3$	$5x^2 + \frac{13}{3}x^3$
$n = 4$	$x + \frac{35}{2}x^2 + 26x^3 + \frac{77}{12}x^4$	$10x^2 + \frac{65}{3}x^3 + \frac{77}{12}x^4$

(102)

Case $\alpha = 3$

${}_r\phi_{n,3}^h(x)$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	x	0
$n = 2$	$x + \frac{7}{2}x^2$	$\frac{7}{2}x^2$
$n = 3$	$x + \frac{21}{2}x^2 + \frac{47}{6}x^3$	$7x^2 + \frac{47}{6}x^3$
$n = 4$	$x + \frac{49}{2}x^2 + 47x^3 + \frac{171}{12}x^4$	$14x^2 + \frac{235}{6}x^3 + \frac{171}{12}x^4$

(103)

Hence "hyperharmonic r -exponential numbers" are defined as

$${}_r\phi_{n,\alpha}^h = \sum_{k=0}^n \left\{ \begin{matrix} n \\ k \end{matrix} \right\}_r H_k^{(\alpha)}. \quad (104)$$

The first few hyperharmonic r -exponential numbers are:
Case $\alpha = 2$

${}_r\phi_{n,2}^h$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	1	0
$n = 2$	$\frac{7}{2}$	$\frac{5}{2}$
$n = 3$	$\frac{77}{6}$	$\frac{28}{3}$
$n = 4$	$\frac{611}{12}$	$\frac{457}{12}$

(105)

Case $\alpha = 3$

${}_r\phi_{n,3}^h$	$r = 1$	$r = 2$
$n = 0$	0	0
$n = 1$	1	0
$n = 2$	$\frac{9}{2}$	$\frac{7}{2}$
$n = 3$	$\frac{58}{3}$	$\frac{89}{6}$
$n = 4$	$\frac{347}{4}$	$\frac{809}{12}$

(106)

Remark 29 We extend the relation (80) as

$${}_rF_{n,\alpha}^h(z) = \int_{0^r}^{\infty} \phi_{n,\alpha}^h(z\lambda) e^{-\lambda} d\lambda.$$

References

- [1] Abramowitz, M and Stegun, I. *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, 9th printing. New York: Dover, p. 824, 1972.
- [2] Bell, E. T. *Exponential polynomials*, Annals of Mathematics, vol. 35, no. 2, pp. 258–277, (1934).
- [3] Bell, E. T. *Exponential numbers*, Amer. Math. Monthly 41, 411-419, (1934).
- [4] Benjamin, A.T., Gaebler D. and Gaebler, R. *A combinatorial approach to hyperharmonic numbers*, Integers: Electron. J. Combin. Number Theory 3 (2003), pp. 1–9 #A15.
- [5] Benjamin, A. T. and Quinn, J. J. *Proofs that Really Count: The Art of Combinatorial Proof*, MAA, 2003.
- [6] Boyadzhiev, Khristo N. *A Series transformation formula and related polynomials*, In. J. Math. Math. Sc. 2005: 23 (2005), 3849-3866.
- [7] Boyadzhiev, Khristo N. *Exponential Polynomials, Stirling Numbers and Evaluation of some Gamma Integrals*, Abstract and Applied Analysis, Volume 2009, Article ID 168672.
- [8] Broder, A. Z. *The r-Stirling numbers*, Discrete Math. 49 (1984), 241-259.
- [9] Carlitz, L. *Weighted Stirling numbers of the first and second kind-I*, The Fibonacci Quarterly, 18 (1980), 147-162.
- [10] Carlitz, L. *Weighted Stirling numbers of the first and second kind-II*, The Fibonacci Quarterly, 18 (1980), 242-257.
- [11] Comtet, L. *Advanced Combinatorics. The Art of Finite and Infinite Expansions, Revised and enlarged edition*, D. Riedel Publishing Co., Dordrecht, 1974.

- [12] Conway J. H. and Guy R. K., *The book of numbers*, New York, Springer-Verlag, 1996.
- [13] Dil, A. and Mezo, I. *A Symmetric Algorithm for Hyperharmonic and Fibonacci Numbers*, Applied Mathematics and Computation 206 (2008), 942–951.
- [14] Dil, A. and Kurt, V. *Investigating Geometric and exponential Polynomials with Euler-Seidel Algorithm*, submitted. Available at <http://arxiv.org/abs/0908.2585>.
- [15] Dil, A. and Kurt, V. *Polynomials Related to Harmonic Numbers and Evaluation of Harmonic Number Series I*, submitted.
- [16] Grunert, J. A. *Über die Summierung der Reihen...*, Journal für die reine und angewandte Mathematik, vol. 25, pp. 240–279, 1843.
- [17] Graham, R. L., Knuth D. E. and Patashnik O., *Concrete Mathematics*, Addison Wesley, 1993.
- [18] Mezo, I. *r-exponential numbers*, arXiv:0909.4417, 2009.
- [19] Mezo, I., Dil, A. *Hyperharmonic series involving Hurwitz zeta function*, Journal of Number Theory, 130, 2, 2010, 360-369.
- [20] Nyul, G. *Ordered r-exponential numbers and r-Eulerian numbers* (in Hungarian), Diophantine and Cryptography Days in Sopron, 11 October 2008, Sopron.
- [21] Riordan, J. *Combinatorial Analysis*, John Wiley, New York, 1958.
- [22] Schwatt, I. J. *An Introduction to the Operations with Series*, Chelsea, New York, 1962.
- [23] Tanny, Stephen M. *On some numbers related to the exponential numbers*, Canadian Mathematical Bulletin, Vol 17, (1974), No 5, 733-738.
- [24] Wilf, Herbert. S. *Generatingfunctionology*, Academic Press, 1993.